

# SOLUTION

---

Ex 8.1: 1, 6, 8, 16, 20

Ex 8.2: 2, 3, 8

Ex 8.3: 1, 4, 6, 9, 10

Ex 8.4 and Ex 8.5: 4, 5, 7, 8, 12

## Ex 8.1: (1)

- Let  $x \in S$  and let  $n$  be the number of conditions (from among  $c_1, c_2, c_3, c_4$ ) satisfied by  $x$ . ( $n = 0$ ): Here  $x$  is counted once in  $N(\bar{c}_2\bar{c}_3\bar{c}_4)$  and once in  $N(\bar{c}_1\bar{c}_2\bar{c}_3\bar{c}_4)$ . ( $n = 1$ ): If  $x$  satisfies  $c_1$  (and not  $c_2, c_3, c_4$ ), then  $x$  is counted once in  $N(\bar{c}_2\bar{c}_3\bar{c}_4)$  and once in  $N(\bar{c}_1\bar{c}_2\bar{c}_3\bar{c}_4)$
- If  $x$  satisfies  $c_i$ , for  $i \neq 1$ , then  $x$  is not counted in any of the three terms in the equation. ( $n = 2, 3, 4$ ): If  $x$  satisfies at least two of the four conditions, then  $x$  is not counted in any of the three terms in the equation.
- The preceding observations show that the two sides of the given equation count the same elements from  $S$ , and this provides a combinational proof for the formula
$$N(\bar{c}_2\bar{c}_3\bar{c}_4) = N(c_1\bar{c}_2\bar{c}_3\bar{c}_4) + N(\bar{c}_1\bar{c}_2\bar{c}_3\bar{c}_4)$$

## Ex 8.1: (6a, 6b)

- $x_1 + x_2 + x_3 + x_4 = 19$

a)  $0 \leq x_i, 1 \leq i \leq 4. \binom{4+19-1}{19} = \binom{22}{19}$

b) For  $1 \leq i \leq 4$ , let  $c_i: x_i \geq 8$ .

$$N(c_i): x_1 + x_2 + x_3 + x_4 = 11: \binom{4+11-1}{11} = \binom{14}{11}, 1 \leq i \leq 4.$$

$$N(c_i c_j): x_1 + x_2 + x_3 + x_4 = 3: \binom{4+3-1}{3} = \binom{14}{3}, 1 \leq i < j \leq 4.$$

$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4) = N - S_1 + S_2 = \binom{22}{19} - 4 \binom{14}{11} + 6 \binom{6}{3}.$$

## Ex 8.1: (6c)

- The number of solutions for  $x_1 + x_2 + x_3 + x_4 = 19$  where  $0 \leq x_1 \leq 5, 0 \leq x_2 \leq 6, 3 \leq x_3 \leq 7, 3 \leq x_4 \leq 8$  equals the number of solutions for  $x_1 + x_2 + x_3 + x_4 = 13$  with  $0 \leq x_1 \leq 5, 0 \leq x_2 \leq 6, 0 \leq x_3 \leq 4, 0 \leq x_4 \leq 5$ . Define the conditions  $c_i, 1 \leq i \leq 4$ , as follows:  $c_1: x_1 \geq 6, c_2: x_2 \geq 7; c_3: x_3 \geq 5; c_4: x_4 \geq 6$ .

$$N = \binom{4+13-1}{13} = \binom{16}{13}.$$

$$N(c_1), N(c_4): x_1 + x_2 + x_3 + x_4 = 7: \binom{4+7-1}{7} = \binom{10}{7}.$$

$$N(c_2): x_1 + x_2 + x_3 + x_4 = 6: \binom{4+6-1}{6} = \binom{9}{6}.$$

$$N(c_3): x_1 + x_2 + x_3 + x_4 = 8: \binom{4+8-1}{8} = \binom{11}{8}.$$

$$N(c_1 c_2) = 1.$$

$$N(c_1 c_3): x_1 + x_2 + x_3 + x_4 = 2: \binom{4+2-1}{2} = \binom{5}{2}.$$

$$N(c_1 c_4): x_1 + x_2 + x_3 + x_4 = 1: \binom{4+1-1}{1} = \binom{4}{1}.$$

$$N(c_2 c_3) = \binom{4}{1}, N(c_2 c_4) = 1, N(c_3 c_4) = \binom{5}{2}.$$

$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4) = \binom{16}{13} - [2\binom{10}{7} + \binom{9}{6} + \binom{11}{8}] + 2[1 + \binom{4}{1} + \binom{5}{2}].$$

## Ex 8.1: (8)

- The number of integer solutions for  $x_1 + x_2 + x_3 + x_4 = 19$ ,  $5 \leq x_i \leq 10$ ,  $1 \leq i \leq 4$ , equals the number of integer solutions for  $y_1 + y_2 + y_3 + y_4 = 39$ ,  $0 \leq y_i \leq 15$ .

- For  $1 \leq i \leq 4$ , let  $c_i: y_i \geq 16$ .

$$N(c_i), 1 \leq i \leq 4: y_1 + y_2 + y_3 + y_4 = 23: \binom{4+23-1}{23} = \binom{26}{23}.$$

$$N(c_i c_j), 1 \leq i < j \leq 4: y_1 + y_2 + y_3 + y_4 = 7: \binom{4+7-1}{7} = \binom{10}{7}.$$

$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4) = \binom{42}{39} - 4 \binom{26}{23} + 6 \binom{10}{7}.$$

## Ex 8.1: (16)

- $10^9 - \binom{3}{1}9^9 + \binom{3}{2}8^9 - \binom{3}{3}7^9.$

## Ex 8.1: (20)

- For  $1 \leq i \leq 7$ , let  $c_i$  denote the situation where the  $i$ -th friend was at lunch with Sharon.

Then  $N(\bar{c}_1 \bar{c}_2 \dots \bar{c}_7) =$

$$84 - \binom{7}{1}35 + \binom{7}{2}16 - \binom{7}{3}8 + \binom{7}{4}4 - \binom{7}{5}2 + \binom{7}{6}1 - \binom{7}{7}0 = 0.$$

Consequently, Sharon always had company at lunch.

## Ex 8.2: (2)

- a) Let  $c_i$  denote the condition that the two A's are together in an arrangement of ARRANGEMENT. Conditions  $c_2, c_3, c_4$  are defined similarly for the two E's, N's, and R's, respectively.

$$N = 2494800.$$

$$\text{For } 1 \leq i \leq 4, N(c_i) = \frac{10!}{(2!)^3} = 453600.$$

$$\text{For } 1 \leq i < j \leq 4, N(c_i c_j) = \frac{9!}{(2!)^2} = 90720.$$

$$N(c_i c_j c_k) = \frac{8!}{2!} = 20160, 1 \leq i < j < k \leq 4.$$

$$N(c_1 c_2 c_3 c_4) = 7! = 5040.$$

$$S_1 = \binom{4}{1} 453600 = 1814400. S_2 = \binom{4}{2} 90720 = 544320.$$

$$S_3 = \binom{4}{3} 20160 = 80640. S_4 = \binom{4}{4} 5040 = 5040.$$

$$(i) E_2 = S_2 - \binom{3}{1} S_3 + \binom{4}{2} S_4 = 332640$$

$$(ii) L_2 = S_2 - \binom{2}{1} S_3 + \binom{3}{1} S_4 = 398160$$

- b) (i)  $E_3 = S_3 - \binom{4}{1} S_4 = 60480.$  (ii)  $L_3 = S_3 - \binom{3}{2} S_4 = 65520.$



## Ex 8.2: (3)

- Let  $c_1$  denote the presence of consecutive E's in the arrangement. Likewise,  $c_2$ ,  $c_3$ ,  $c_4$ , and  $c_5$  are defined for consecutive N's, O's, R's, and S's, respectively.

a) 
$$N = \frac{14!}{(2!)^5}$$
$$N(c_1) = \frac{13!}{(2!)^4}; S_1 = \binom{5}{1} \left( \frac{13!}{(2!)^4} \right).$$
$$N(c_1 c_2) = \frac{12!}{(2!)^3}; S_2 = \binom{5}{2} \left( \frac{12!}{(2!)^3} \right).$$
$$N(c_1 c_2 c_3) = \frac{11!}{(2!)^2}; S_3 = \binom{5}{3} \left( \frac{11!}{(2!)^2} \right).$$
$$N(c_1 c_2 c_3 c_4) = \frac{10!}{(2!)^1}; S_4 = \binom{5}{4} \left( \frac{10!}{(2!)^1} \right).$$
$$N(c_1 c_2 c_3 c_4 c_5) = 9! = S_5.$$
$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4 \bar{c}_5) = 1,286,046,720$$

b) 
$$E_2 = S_2 - \binom{3}{1} S_3 + \binom{4}{2} S_4 - \binom{5}{3} S_5 = 350,179,200$$

c) 
$$L_3 = S_3 - \binom{3}{2} S_4 + \binom{4}{2} S_5 = 74,753,280$$

## Ex 8.2: (8)

b)  $E_{t-1} = S_{t-1} - tS_t; L_{t-1} = L_t + E_{t-1}$

c)  $L_{t-1} = L_t + E_{t-1} = S_t + S_{t-1} - tS_t = S_{t-1} - (t-1)S_t = S_{t-1} - \binom{t-1}{t-2}S_t$

d)  $L_m = L_{m+1} + E_m$

e)  $L_t = S_t$

$$L_{t-1} = S_{t-1} - \binom{t-1}{t-2}S_t$$

Assume  $L_{k+1} = S_{k+1} - \binom{k+1}{k}S_{k+2} + \binom{k+2}{k}S_{k+3} - \dots + (-1)^{t-k-1}\binom{t-1}{k}S_t$

$$L_k = L_{k+1} + E_k =$$

$$\left[ S_{k+1} - \binom{k+1}{k}S_{k+2} + \binom{k+2}{k}S_{k+3} - \dots + (-1)^{t-k-1}\binom{t-1}{k}S_t \right]$$

$$+ \left[ S_k - \binom{k+1}{1}S_{k+1} + \binom{k+1}{2}S_{k+2} - \dots + (-1)^{t-k}\binom{t}{t-k}S_t \right].$$

For  $1 \leq r \leq t-k$ , the coefficient of  $S_{k+r}$  is  $(-1)^{r-1}\binom{k+r-1}{k} + (-1)^r\binom{k+r}{r} = (-1)^r\binom{k+r-1}{k-1}$ .

Consequently,  $L_k = S_k - \binom{k}{k-1}S_{k+1} + \binom{k+1}{k-1}S_{k+2} - \dots + (-1)^{t-k}\binom{t-1}{k-1}S_t$ .

## Ex 8.3: (1)

- For  $1 \leq i \leq 5$  let  $c_i$  be the condition that  $2i$  is in position  $2i$ .

$$N = 10!; N(c_i) = 9!; N(c_i c_j) = 8!, 1 \leq i < j \leq 5; \dots;$$

$$N(c_1 c_2 c_3 c_4 c_5) = 5!.$$

$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4 \bar{c}_5) = 10! - \binom{5}{1}9! + \binom{5}{2}8! - \binom{5}{3}7! + \binom{5}{4}6! - \binom{5}{5}5!$$

## Ex 8.3: (4)

- There are  $7! = 5040$  permutations for  $1,2,3,4,5,6,7$ .

Among these there are  $7! \left[ 1 - 1 + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \frac{1}{5!} + \frac{1}{6!} - \frac{1}{7!} \right] =$   
1851 derangements.

Consequently, we have  $5040 - 1854 = 3186$  permutations of  $1,2,3,4,5,6,7$  that are not derangements.

## Ex 8.3: (6)

- a) There are  $(d_4)^2 = 9^2 = 81$  such derangements.
- b) In this case we get  $(4!)^2 = 24^2 = 576$  derangements.

## Ex 8.3: (9)

- $(10!)d_{10} = (10!)^2(e^{-1})$

## Ex 8.3: (10)

a) (i)  $\frac{d_n}{n!}$ . (ii)  $\frac{n(d_{n-1})}{n!}$ . (iii)  $1 - \frac{d_n}{n!}$ . (iv)  $\frac{\binom{n}{r}d_{n-r}}{n!}$ .

b) (i)  $e^{-1}$ . (ii)  $e^{-1}$ . (iii)  $1 - e^{-1}$ . (iv)  $\left(\frac{1}{r!}\right) e^{-1}$ .

## Ex 8.4 and Ex 8.5: (4)

- $r(C_1, x) = 1 + 4x + 3x^2 = r(C_2, x)$



## Ex 8.4 and Ex 8.5: (5)

- a) (i)  $(1 + 2x)^3$   
(ii)  $1 + 8x + 12x^2 + 4x^3$   
(iii)  $1 + 9x + 25x^2 + 21x^3$   
(iv)  $1 + 8x + 16x^2 + 7x^3$
- b) If the board  $C$  consists of  $n$  steps, and each step has  $k$  blocks, then  $r(C, x) = (1 + kx)^n$ .

## Ex 8.4 and Ex 8.5: (7)

- $r(C, x) = (1 + 4x + 3x^2)(1 + 4x + 2x^2)$   
 $= 1 + 8x + 21x^2 + 20x^3 + 6x^4.$

For  $1 \leq i \leq 5$  let  $c_i$  be the condition that an assignment is made with person ( $i$ ) assigned to a language he or she wishes to avoid.

$$N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4 \bar{c}_5) = 5! - 8 \times 4! + 21 \times 3! - 20 \times 2! + 6 \times 1! = 20.$$

	Java	C++	VHDL	SQL
(1) Jeanne				
(2) Charles				
(3) Todd				
(4) Paul				
(5) Sandra				

## Ex 8.4 and Ex 8.5: (8)

- The factor  $6!$  is needed because we are counting ordered sequences.

## Ex 8.4 and Ex 8.5: (12)

- Consider the chessboard  $C$  of shaded squares.

Here  $r(C, x) = 1 + 8x + 20x^2 + 17x^3 + 4x^4$ . For any one-to-one function  $f: A \rightarrow B$ , let  $c_1, c_2, c_3, c_4$  denote the conditions:

$$c_1: f(1) = v \text{ or } w \quad c_3: f(3) = x$$

$$c_2: f(2) = u \text{ or } w \quad c_4: f(4) = v, x, \text{ or } y$$

The answer to this problem is . So there are 146 one-to-one functions  $f: A \rightarrow B$  where

$$f(1) \neq v, w$$

$$f(3) \neq x$$

$$f(2) \neq u, w$$

$$f(4) \neq v, x, y.$$

	u	v	w	x	y	z
1		■	■			
2	■		■			
3				■		
4		■		■	■	
5						